

ເອກສາຮ້ອງອີງ

- ຈິນທານ ກຍາຊຽວມັນ ນິກາ ຈັນທະວີສົມພາຍ ແລະ ວັນການ ຕົວດັນທັກທີ 2539 ສື່ວັນດີຂອງແມ່ລວມບ້າວ Orseolia oryzae (Wood-Mason) ໃນປະເທດໄທ. ມັນຕີ 604-650. ໃນ: ວາງຈານປະຈຳປີ 2539 ດັວກປະຫຼວມສັນນາກາງວິທະການຂອງແມ່ລວມແລະຄັດຮູ້ທີ່ກໍານົດວິທະການ ດັວກປະຫຼວມ ກຽມວິທະການເກຍເທດ ກຽມເທດ ຖະໜາກ ດັວກປະຫຼວມສັນນາໃຈປົງບົດກົດຮອງກຸ່ມເກຍເທດກຣ. ອ. ເມ່ວະນະກຳ ຈ. ດັກ 7 ມັນຕີ ດໍາເນີນ ກາລະດີ ສັນສົນທີ່ ຈຳດັດ ແສງວິວາ ສູງຍິງຄໍ ກົນກວຽຮຣ ສົງຈານ ປົມືດາ ບຸນູສູທິ່ ແລະສົມພານ ນັນ ພະເສນ 2552 ສາຮແກນມ່າໄໂໄໂໄ ໄສ່ນອດແລະແອນໄກ ໄໃຫຍນີນໃນຄວາມຫລາກຫາຍາງ ພັນຖຸກຽມຂອງໜ້າວ່າເກີນເມື່ອຈີ່ໄທຢະແກນເປັນໄປໄດ້ໃນກາງຮ້າງພັນຖຸໃຫ້ມ່ເພື່ອເພີ່ມຄຸນຄໍາ ຄວາມເປັນໄໝ້າວ່າເພື່ອສຸຂະພາບສາບັນວິຊີແລະພັດນາວິທະຫາສາສົກຮ່ວມເກົດໄນ ໂລື້ ມາກວິທະກູ້ເກີນເປັນໄປໃໝ່ ພ.ສ. 2552 138 ມັນຕີ
- ພອນນີ້ ຖຸການຈົກ 2549 ຄວາມຫລາກຫາຍາງພັນຖຸກຽມຂອງໜ້າວ່າເກີນເມື່ອຈີ່ໄທເໜີນອອງ ວິທະນີນີ້ວິທະຫາສາດຕຽມຫານບັດທິດ. ມາກວິທະກູ້ເກີນເປັນໄປໃໝ່ ເຊິ່ງໃໝ່ 94 ມັນຕີ
- ຮັດຕິຫາ ທະຫາພົກ 2549 ຄວາມຫລາກຫາຍາງສາບັນຫຼຸຂອງປະຈາກແມ່ລວມບ້າວິກາເກີນຂອງປະເທດໄທ ວິທະນີນີ້ວິທະຫາສາດຕຽມຫານບັດທິດ ມາກວິທະກູ້ເກີນເປັນໄປໃໝ່ ເຊິ່ງໃໝ່ 71 ມັນຕີ ສັນສົນທີ່ ຈຳດັດ ນິສີ ອື່ນແໜ້ນ ຜາການທີ່ເກີນໄດ້ໂທໄບໂລດ໌ ພຣົມຖຸທີ່ແລະ ສັກທີ່ຂຶ້ນ ລອດເກົ່ວ 2558 ໂຄງກາງວິຊີ ເຊິ່ງບູຮັມກາເຮົາເພື່ອສ່ວນຮ້າງປະສົງວິທະການກາງການຄຸດແລະກາງການຄຸດຂອງໜ້າວ່ານີ້ທີ່ສູງ ໂຄງກາງຍ່ອຍທີ່ 4: ກາງວິຊີປ່ຽນປ່ຽນປ່ຽນຖຸ້ຫາວ່ານີ້ທີ່ສູງທີ່ກຳນົດການດ້ວຍແມ່ລວມບ້າວິກາເກີນເປັນໄປໃໝ່ ຄຸນກາຫົວໜ້າກາງໂກ່ງນາການ ຮ້າງຈານຈົບັນສົມບູຮັມ໌ ສາບັນວິຊີແລະພັດນາເກີນເປັນໄປໃໝ່ (ອົງກໍາການມາຫານ) ປະຈຳປິ່ງປະປະມາພ. ພ.ສ. 2558 73 ມັນຕີ
- ສັນສົນທີ່ ຈຳດັດ ນິສີ ອື່ນແໜ້ນ ຜາການທີ່ເກີນໄດ້ໂທໄບໂລດ໌ ພຣົມຖຸທີ່ແລະ ສັກທີ່ຂຶ້ນ ລອດເກົ່ວ 2557 ໂຄງກາງວິຊີ ເຊິ່ງບູຮັມກາເຮົາເພື່ອສ່ວນຮ້າງປະສົງວິທະການວິທະການຮອມອິດແລະກາງການຄຸດຂອງໜ້າວ່ານີ້ທີ່ສູງ ໂຄງກາງຍ່ອຍທີ່ 4: ກາງໄໝ້ປະໄວ ໂອນແລະປັບປຸງພັນຖຸ້ຫາວ່າອ່ອງເຖິງດຳກຳນົດຮູ້ຫຼຸມນັນນີ້ທີ່ສູງ ຮ້າງຈານຈົບັນສົມບູຮັມ໌ ສາບັນວິຊີແລະພັດນາເກີນເປັນໄປໃໝ່ (ອົງກໍາການມາຫານ) ປະຈຳປິ່ງປະປະມາພ. ພ.ສ. 2557 65 ມັນຕີ
- Black, R., Lindsay, H., Bhutta, Z., Caulfield, L. and de Onnis, M. 2008. Maternal and child under-nutrition: global and regional exposures and health consequences. *Lancet.* 371: 243-260.
- Boonsit, P., Pongpiachan, P., Julsrigival, S. and Karladee, D. 2010. Gamma oryzanol content in glutinous purple rice landrace varieties. *CMU. J. Nat. Sci.* 9: 151-157.
- Brown, A.H.D. 1978. Isozymes, plant population genetics structure and genetic conservation. *Theor. Appl. Genet.* 52: 145–157

- Daiponmak, W., Theerakulpisut, P., Thanonkao, P., Vanavichit, A. and Prathepha, P. 2010. Changes of anthocyanin cyanidin-3-glucoside content and antioxidant activity in Thai rice varieties under salinity stress. *ScienceAsia*. 36: 286-291.
- Graham, R.D., Senadhira, D., Beebe, S.E., Iglesias, C. and Ortiz-Monasterio, I. 1999. Breeding for micronutrient density in edible portions of staple food crops: conventional approaches. *Field Crops Research*. 60:57-80
- Harlan, J. 1992. Crop and Man. 2nd Ed. Am Soc Agron, Madison, Wisconsin. 284p.
- Hettiarachchi, M., Hilmers, D. C., Liyanage, C. and Abrams, S.A. 2004. Na(2)EDTA enhances the absorption of iron and zinc from fortified rice flour in Sri Lankan children. *J. Nutr.* 134: 3031-3036.
- Hidaka, T., Vungsilburt, P. and Kadkao, S. 1974. Studies on Ecology and Control of the Rice Gall Midge in Thailand. Technical Bulletin No.6, TARC. 113 pp.
- International Food Policy Research Institute 1999. World Food Prospects: Critical Issue for the Early Twenty- first Century.
- Juliano, B. O. 1993. Rice in human nutrition "Prepared in collaboration with FAO". Rome: Rome: Food and Agriculture Organization of the United Nations.
- Lee, J.H. 2010. Identification and quantification of anthocyanins from the grains of black rice (*Oryza sativa* L.) varieties. *Food Sci. Biotechnol.* 19: 391-397.
- Mahatheeranont, S., Keawsa-ard, S. and Dumri, K. 2001. Quantification of the rice aroma compound, 2-Acetyl-1-pyrroline, in uncooked Khao Dawk Mali 105 brown rice. *J. J. Agric. Food Chem.* 49: 773-779.
- Muntana, N., Prasong, S. 2010. Study on total phenolic contents and their antioxidant activities of Thai white, red, and black rice bran extracts. *Pakistan J. Biol. Sci.* 13: 170-4.
- Okai, Y., Higashi-Okai, K. 2006. Radical-scavenging activity of hot water extract of Japanese rice bran— association with phenolic acids. *J. UOEH.* 28:1-12.
- Upkaew, P., Pusadee, T., Sirabanchongkran, A., Rerkasem, K., Jamjod, S., Rerkasem, B. 2011. Complexity and adaptability of a traditional agricultural system: case study of a gall midge resistant rice landrace from northern Thailand. *Genet. Resour. Crop Ev.* 58:361-372.
- Phattarakul, N. 2008. Genotypic variation in tolerance to acid soil in local upland rice varieties. Ph.D. Thesis, Graduate School. Chiang Mai University, Chiang Mai, Thailand.

- Pintasen, S., Prom-u-thai, C., Jamjod, S., Yimyam, N. and Rerkasem, B. 2007. Variation of grain iron content in a local upland rice germplasm from the village of Huai Tee Cha in northern Thailand. *Euphytica*. 158:27-34.
- Pintasen, S., Prom-u-thai, C., Jamjod, S., Yimyam, N. and Rerkasem, B. 2007. Variation of grain iron content in a local upland rice germplasm from the village of Huai Tee Cha in northern Thailand. *Euphytica*. 158: 27-34.
- Prom-u-thai, C. 2003. Iron (Fe) in Rice grain. Ph.D. Thesis, Graduate School, Chiang Mai University, Chiang Mai, Thailand.
- Prom-u-thai, C. and Rerkasem, B. 2001. Grain iron concentration in Thai rice germplasm. Plant nutrition–Food security and sustainability of agro-ecosystems. 350-351
- Prom-u-thai, C., Fukai, S., Godwin, I., Huang, L. 2007. Genotypic variation of iron partitioning in rice grain. *J. Sci. Food Agric.* 87: 2049-2054.
- Prom-u-thai, C., Sanchai, C., Rerkasem, B., Jamjod, S., Fukai, S., Godwin, I.D., Huang, L. 2007. Grain morphology variability and its effect to degree of milling and Fe loss in rice. *Cereal Chem.* 84: 384-388.
- Pusadee, T., Jamjod, S., Chiang, Y., Rerkasem, B. and Schaal, BA. 2009. Genetic structure and isolation by distance in a landrace of Thai rice. *Proc. Nat. Acad. Sci. USA*. 106: 13880-13885.
- Rosado, J. 2003. Zinc and Copper: Proposed Fortification Levels and Recommended Zinc Compounds. *J. Nutr.* 133: 2985S-2589S.
- Ryu, S.N., Park, S.Z., and Ho, C.T. 1998. High performance liquid chromatographic determination of anthocyanin pigments in some varieties of black rice. *J. Food Drug Anal.* 6: 729-736.
- Sardesai, N., Rajyashri, K.R., Behura, S.K., Nair, S. and Mohan, M. 2001. Genetic, physiological and molecular interactions of rice and its major dipteran pest, gall midge. *Plant Cell, Tissue and Organ Culture*. 64: 115-131.
- Suwannalert, P. and Rattanachithwat, S.,2011. High level of phytophenolics and antioxidant activities in *Oryza sativa* - unpolisded Thai rice strain of LeumPhua, *Trop J. Pharmaceut Res.* 10(4): 431-436.
- Tananuwong, K. and Tewaruth, W.,2010. Extraction and application of antioxidants from black glutinous rice.*Food Sci Tech.* 43: 476-481.

- Thongphak, D., Attathom, T. and Tayathum, C. 1999. Determinaton of the genetic relatedness of the rice gall midge, *Orseolia oryzae* in Thailand using RAPD-PCR marker. Thai Journal of Agricultural Science. 32: 409-421.
- Zarcinas, B. A., Cartwright, B. and Spouncer, L.R. 1987. Nitric acid digestion and multi-element analysis of plant material by inductively coupled plasma spectrometry. Communication of Soil Science and Plant Analysis. 18: 131-146.

